

Monaco, 22 March 2007 - Georges MOUTON MD

MULTIPLE SCLEROSIS: A FUNCTIONAL APPROACH

The Lancet has published several articles and letters between 1976 and 1978 reporting small bowel abnormalities in multiple sclerosis including increased inflammatory-cell infiltration [1] and loss of villous architecture [2, 3]. Several authors, including in *The American Journal of Gastroenterology*, linked cases of multiple sclerosis with intestinal malabsorption [2-4].

We still do not well understand the mechanisms underlying the initiation and progression of multiple sclerosis, but it has an autoimmune origin [5]. However, "unlike autoimmune conditions such as rheumatoid arthritis, systemic lupus erythematosus or myasthenia gravis, multiple sclerosis has no specific immunological marker" [6], despite the fact that the presence of at least two different antibodies - anti-myelin oligodendrocyte glycoprotein (MOG) and anti-myelin basic protein (MBP) - appears highly predictive of typical clinical events that establish the diagnosis of clinically definite multiple sclerosis [7].

Recently, more evidence accumulates about the link between autoimmune diseases and an abnormal intestinal permeability, a condition often named "*leaky gut syndrome*". An article published in *Science* (March 2005) suggests that a leaky gut barrier allows antigens to enter the lamina propria, leading to the disruption of the immune system [8]. Another paper that appeared soon after in *Acta Paediatrica* (April 2005) explained how leaky intestines could trigger an abnormal antigen exposure, itself enabling the cross-recognition of molecular structures shared between self and nonself, a phenomenon called "molecular mimicry" [9].

Fasano, the world-leading expert on celiac disease, claims that "the loss of the protective function of mucosal barriers (mainly the gastrointestinal and lung mucosa) is necessary for autoimmunity to develop" [10]. He even suggests, "the autoimmune process can be arrested if the interplay between genes and environmental triggers is prevented by re-establishing intestinal barrier function" [10]. Actually, Crohn's disease as the other inflammatory bowel diseases corresponds to an autoimmune condition characterized by increased intestinal permeability [11]. Interestingly, "multiple sclerosis (MS) patients may have an increased risk of co-acquisition of Crohn's disease" and "a subgroup (5 among 20 patients enrolled in the study) of MS patients has increased intestinal permeability" [12].

Taking into account that "alcohol abuse impairs the function of the intestinal barrier" [13], and that ethanol is known to increase intestinal permeability [14], we underline how MS patients are well known to be at increased risk from alcohol excessive consumption [15]. Consequently, several paths may lead to the gut metabolism abnormalities seen in MS [16].

"Intestinal permeability is characteristically elevated in untreated celiac disease" [17] as well as "intestinal permeability is altered in children with cow's milk allergy" [18]. In fact, impaired intestinal permeability seems to be present in all subjects with adverse reactions to food [19] and breast-feeding could favourably influence intestinal permeability [19]. Not surprisingly, dairy-free diet [20] and gluten-free diet [21] could show beneficial to MS patients, and they "were less likely than controls to have been breast fed for a prolonged period of time, another point [being] added to the long list of the benefits of prolonged breast feeding" [22]. We spot similar beneficial effects towards autistic patients [23, 24], while these patients also suffer from an abnormal intestinal permeability [25].

1. Lange, L.S. and M. Shiner, *Small-bowel abnormalities in multiple sclerosis*. Lancet, 1976. **2**(7999): p. 1319-22.
2. Cook, A.W., et al., *Multiple sclerosis and malabsorption*. Lancet, 1978. **1**(8078): p. 1366.
3. Fantelli, F.J., H. Mitsumoto, and B.A. Sebek, *Multiple sclerosis and malabsorption*. Lancet, 1978. **1**(8072): p. 1039-40.
4. Gupta, J.K., et al., *Multiple sclerosis and malabsorption*. Am J Gastroenterol, 1977. **68**(6): p. 560-5.
5. Olson, J.K., et al., *A virus-induced molecular mimicry model of multiple sclerosis*. J Clin Invest, 2001. **108**(2): p. 311-8.
6. Chaudhuri, A. and P.O. Behan, *Multiple sclerosis: looking beyond autoimmunity*. J R Soc Med, 2005. **98**(7): p. 303-6.
7. Antel, J.P. and A. Bar-Or, *Do myelin-directed antibodies predict multiple sclerosis?* N Engl J Med, 2003. **349**(2): p. 107-9.
8. Macdonald, T.T. and G. Monteleone, *Immunity, inflammation, and allergy in the gut*. Science, 2005. **307**(5717): p. 1920-5.
9. Liu, Z., N. Li, and J. Neu, *Tight junctions, leaky intestines, and pediatric diseases*. Acta Paediatr, 2005. **94**(4): p. 386-93.
10. Fasano, A. and T. Shea-Donohue, *Mechanisms of disease: the role of intestinal barrier function in the pathogenesis of gastrointestinal autoimmune diseases*. Nat Clin Pract Gastroenterol Hepatol, 2005. **2**(9): p. 416-22.
11. Welcker, K., et al., *Increased intestinal permeability in patients with inflammatory bowel disease*. Eur J Med Res, 2004. **9**(10): p. 456-60.
12. Yacyshyn, B., et al., *Multiple sclerosis patients have peripheral blood CD45RO+ B cells and increased intestinal permeability*. Dig Dis Sci, 1996. **41**(12): p. 2493-8.
13. Parlesak, A., et al., *Increased intestinal permeability to macromolecules and endotoxemia in patients with chronic alcohol abuse in different stages of alcohol-induced liver disease*. J Hepatol, 2000. **32**(5): p. 742-7.
14. Kiefer, D. and L. Ali-Akbarian, *A brief evidence-based review of two gastrointestinal illnesses: irritable bowel and leaky gut syndromes*. Altern Ther Health Med, 2004. **10**(3): p. 22-30; quiz 31, 92.
15. Quesnel, S. and A. Feinstein, *Multiple sclerosis and alcohol: a study of problem drinking*. Mult Scler, 2004. **10**(2): p. 197-201.
16. Westall, F.C., *Molecular mimicry revisited: gut bacteria and multiple sclerosis*. J Clin Microbiol, 2006. **44**(6): p. 2099-104.
17. Johnston, S.D., M. Smye, and R.P. Watson, *Intestinal permeability tests in coeliac disease*. Clin Lab, 2001. **47**(3-4): p. 143-50.
18. Kalach, N., et al., *Intestinal permeability in children: variation with age and reliability in the diagnosis of cow's milk allergy*. Acta Paediatr, 2001. **90**(5): p. 499-504.
19. Ventura, M.T., et al., *Intestinal permeability in patients with adverse reactions to food*. Dig Liver Dis, 2006. **38**(10): p. 732-6.
20. Malosse, D. and H. Perron, *Correlation analysis between bovine populations, other farm animals, house pets, and multiple sclerosis prevalence*. Neuroepidemiology, 1993. **12**(1): p. 15-27.
21. Reichelt, K.L. and D. Jensen, *IgA antibodies against gliadin and gluten in multiple sclerosis*. Acta Neurol Scand, 2004. **110**(4): p. 239-41.
22. Pisacane, A., et al., *Breast feeding and multiple sclerosis*. Bmj, 1994. **308**(6941): p. 1411-2.
23. Knivsber, A.M., K.L. Reichelt, and M. Nodland, *Reports on dietary intervention in autistic disorders*. Nutr Neurosci, 2001. **4**(1): p. 25-37.
24. Knivsberg, A.M., et al., *A randomised, controlled study of dietary intervention in autistic syndromes*. Nutr Neurosci, 2002. **5**(4): p. 251-61.
25. D'Eufemia, P., et al., *Abnormal intestinal permeability in children with autism*. Acta Paediatr, 1996. **85**(9): p. 1076-9.